

Research Note

New Host and Distribution Records for Coccidia (Apicomplexa: Eimeriidae) from North American Lizards (Reptilia: Sauria)

CHRIS T. McALLISTER,¹ STEVE J. UPTON,² AND STANLEY E. TRAUTH³

¹ Renal-Metabolic Lab (151-G), Department of Veterans Affairs Medical Center,
4500 S. Lancaster Road, Dallas, Texas 75216,

² Division of Biology, Ackert Hall, Kansas State University, Manhattan, Kansas 66506 and

³ Department of Biological Sciences, Arkansas State University, State University, Arkansas 72467

ABSTRACT: Four hundred twenty-three lizards, representing 33 species within 7 families (Anguidae, Crotophytidae, Gekkonidae, Phrynosomatidae, Polychridae, Scincidae, Teiidae) were examined for coccidia. Eighty-two (19%) lizards harbored coccidia. Eighty-four percent of the infected lizards were gekkonids (69/139 or 50%) while 6 non-gekkonid families (13/283 or 5%) accounted for 16% of infected lizards. New host records are reported for *Isoospora scinci* Upton, McAllister, and Trauth, 1991, from the broadhead skink, *Eumeces laticeps*, and *Eimeria sceloporis* Bovee and Telford, 1965, from the crevice spiny lizard, *Sceloporus poinsettii*. In addition, this represents the first report of *E. sceloporis* from Texas.

KEY WORDS: Apicomplexa, Coccidia, Eimeriidae, *Isoospora scinci*, *Eimeria sceloporis*, lizards, Gekkonidae, survey, prevalence, Reptilia, Sauria.

Between March 1986 and December 1993, we conducted surveys on protozoan parasites of select herpetofauna of the southcentral and southwestern United States and Mexico and found several species of lizards harboring coccidia. Further examination revealed 6 previously undescribed coccidia, which we have since described as new (McAllister et al., 1988, 1990, 1991a; Upton et al., 1988, 1991), or either redescribed (McAllister and Upton, 1989) or provided new host records (McAllister et al., 1991b). Herein, we add a new locality record and 2 additional host records and summarize our data on the coccidia of 7 families of North American lizards.

Four hundred twenty-three saurians (see Table 1) were collected by hand, noose, or shot with .22 caliber rat shot from 16 counties or parishes in Arkansas, 24 in Texas, 2 in Louisiana, and 1 each in New Mexico and Oklahoma, and from the state of Veracruz, Mexico. Specimens were returned to the laboratory and intestinal contents and feces were examined for coccidia following previously published methods (Upton et al., 1988). Except for lizards collected in Mexico, which were released, host voucher specimens are

deposited in the Arkansas State University Museum of Zoology (ASUMZ) and the Carnegie Museum of Natural History (CM).

Of the 423 individual lizards, representing 33 species within 7 families, 82 (19%) were found to harbor 9 species of coccidians (Table 1). Eighty-four percent of these were gekkonids as 69 of 139 (50%) had 4 species of eimerians. The remaining 16% of the infected lizards were in 4 families as 3 of 131 (2%) of the phrynosomatids had a single eimerian, 2 of 14 (14%) of the polychrids had a caryosporan, 7 of 59 (12%) of the scincids had an eimerian and an isosporan, and 1 of 58 (2%) of the teiids had an eimerian. Three anguids and 18 crotaphytids were negative. There is a significant difference in prevalence of coccidia among gekkonids and non-gekkonids ($\chi^2 = 118.0$, 1 df, $P < 0.00001$). If gekkonids are excluded from the data set, only 13 of 283 (5%) of the lizards were infected.

A new host record was documented for *Isoospora scinci* Upton, McAllister, and Trauth, 1991, in the broadhead skink, *Eumeces laticeps* (Schneider, 1801). The infected juvenile male skink (ASUMZ 19148, snout–vent length = 70 mm) was collected on 18 August 1993 in Independence County, Arkansas, 12.1 km NW Possum Grape at Gold Mine Springs. Three additional Arkansas *E. laticeps*, 2 from Desha County and 1 from Searcy County were negative. *Isoospora scinci* was originally described from five-lined skinks, *Eumeces fasciatus* (Linnaeus, 1758) from Van Buren and Woodruff counties, Arkansas (Upton et al., 1991). These sites are 111 km west and 54 km southeast of the new locale, respectively. This finding was not surprising given that these skinks are broadly sympatric throughout their range (Conant and Collins, 1991).

Sporulated oocysts of an eimerian matching the description of *Eimeria sceloporis* Bovee and

Table 1. Lizards surveyed and the coccidian species collected.

Lizard family/species*	Locality	Prevalence†	Coccidian
Anguidae			
<i>Ophisaurus attenuatus</i>	Arkansas	0/2 (0%)	—
	Texas	0/1 (0%)	—
Crotaphytidae			
<i>Crotaphytus collaris</i>	Texas	0/17 (0%)	—
<i>C. reticulatus</i>	Texas	0/1 (0%)	—
Gekkonidae			
<i>Cyrtopodion scabrum</i>	Texas	8/20 (40%)	<i>Eimeria lineri</i>
<i>Hemidactylus frenatus</i>	Texas	7/12 (58%)	<i>E. dixonii</i>
<i>H. mabouia</i>	Mexico	1/1 (100%)	<i>E. boveroi</i>
<i>H. turcicus</i>	Louisiana	3/7 (43%)	<i>E. lineri</i>
	Texas	36/99 (36%)	<i>E. lineri</i>
	Texas	14/99 (14%)	<i>E. turcicus</i>
Phrynosomatidae			
<i>Cophosaurus texanus</i>	Texas	0/65 (0%)	—
<i>Holbrookia lacerata</i>	Texas	0/1 (0%)	—
<i>H. propinqua</i>	Texas	0/1 (0%)	—
<i>Phrynosoma cornutum</i>	Texas	0/6 (0%)	—
<i>Sceloporus cyanogenys</i>	Texas	0/1 (0%)	—
<i>S. olivaceus</i>	Texas	0/24 (0%)	—
<i>S. poinsettii</i>	Texas	1/4 (0%)	<i>E. sceloporis</i>
<i>S. undulatus</i>	Arkansas	0/9 (0%)	—
	New Mexico	0/1 (0%)	—
	Texas	0/1 (0%)	—
<i>S. variabilis</i>	Mexico	1/1 (100%)	<i>E. sceloporis</i>
	Texas	0/1 (0%)	—
<i>Urosaurus ornatus</i>	Texas	0/7 (0%)	—
<i>Uta stansburiana</i>	Texas	0/8 (0%)	—
Polychridae			
<i>Anolis carolinensis</i>	Arkansas	0/4 (0%)	—
	Louisiana	2/10 (20%)	<i>Caryospora ernsti</i>
Scincidae			
<i>Eumeces anthracinus</i>	Arkansas	0/2 (0%)	—
<i>E. fasciatus</i>	Arkansas	3/14 (21%)	<i>Eimeria fasciatus</i>
	Arkansas	3/14 (21%)	<i>Isospora scinci</i>
<i>E. laticeps</i>	Arkansas	1/4 (25%)	<i>I. scinci</i>
<i>E. obsoletus</i>	Texas	0/2 (0%)	—
<i>E. septentrionalis</i>	Texas	0/3 (0%)	—
<i>E. tetragrammus</i>	Texas	0/8 (0%)	—
<i>Scincella laterale</i>	Arkansas	0/15 (0%)	—
	Oklahoma	0/5 (0%)	—
	Texas	0/6 (0%)	—
Teiidae			
<i>Cnemidophorus gularis</i>	Texas	0/22 (0%)	—
<i>C. inornatus</i>	Texas	0/2 (0%)	—
<i>C. laredoensis</i>	Texas	0/2 (0%)	—
<i>C. septemvittatus</i>	Texas	0/1 (0%)	—
<i>C. sexlineatus</i>	Arkansas	1/28 (4%)	<i>E. sexlineatus</i>
	Texas	0/2 (0%)	—
<i>C. tessellatus</i>	New Mexico	0/1 (0%)	—

* Family nomenclature for iguanian lizards follows Frost and Etheridge (1989) and current scientific names follow Collins (1990).

† Number infected/number examined (percent).

Table 2. Summary of hosts, localities, and prevalence of *Eimeria sceloporis*.

Host	Locality	Prevalence	Reference
<i>Sceloporus clarkii</i>	Mexico	4/4 (100%)	Bovee and Telford, 1965a
<i>S. jarrovi</i>	Arizona	11/38 (31%)	Mitschler et al., 1993
<i>S. magister</i>	California	1/1 (100%)	Bovee and Telford, 1965a
		1/52 (2%)	Telford, 1970
<i>S. poinsettii</i>	Texas	1/4 (25%)	This paper
<i>S. occidentalis</i>	California	?	Bovee and Telford, 1965a
	Washington	149/178 (84%)	Clark and Colwell, 1973
<i>S. variabilis</i>	Mexico	2/2 (100%)	McAllister and Upton, 1989

Telford, 1965, were found in the feces of a gravid adult female (ASUMZ 19021, SVL = 119 mm) crevice spiny lizard, *Sceloporus poinsettii* Baird and Girard, 1852. The host was collected on 28 May 1993 from Llano County, Texas, 3.2 km N Enchanted Rock State Park, off FM 965. *Eimeria sceloporis* was originally described from Clark's spiny lizards, *Sceloporus clarkii* Baird and Girard, 1852, western fence lizards, *Sceloporus occidentalis* Baird and Girard, 1852, and desert spiny lizards, *Sceloporus magister* Hallowell, 1854 (Bovee and Telford, 1965a). This coccidian is apparently genus specific for it was reported from 6 species of sceloporine lizards of the southwestern and Pacific northwestern United States and Mexico (Table 2). In addition, Mitschler et al. (1993) recently provided a redescription of *E. sceloporis* from Yarrow's spiny lizard, *Sceloporus jarrovi* Cope, 1875.

It is somewhat difficult to explain the disparity in prevalence of coccidia among the lizard families. However, the number of previous reports of coccidia from saurian hosts (other than gekkonids) in North America suggests a low prevalence of infection. Indeed, Matuschka and Bannert (1987) and Matuschka (1989) collectively list 30 species of *Eimeria* and 19 species of *Isospora* from lizards, excluding gekkonids. Of the species listed, only 20% of the eimerians and 11% of the isosporans are known from North America (6 from phrynosomatids, 2 from xantusids, and 1 from a scincid). In addition, all of the lizards that have been reported to serve as hosts for coccidia were collected from humid regions of the extreme western or Gulf Coastal area of the United States, in California, Louisiana, and Washington (Bovee and Telford, 1965a, 1965b; Bovee, 1966, 1969; Clark, 1970; Pellérdy, 1974; Atkinson and Ayala, 1987).

It is possible that many of the lizards from study sites did not serve as suitable hosts for coccidia because of defecation habits. Most of

the lizards were observed to bask upon limestone outcroppings, logs, and miscellaneous debris. Feces deposited on the substrate were desiccated within minutes. During periods of increased lizard activity, typical substrate temperatures can range from 35 to 45°C. If these conditions predominate over the period of lizard activity, the upper high temperatures would not represent ideal conditions for coccidia to develop or remain viable. The process of oocyst sporulation is temperature dependent (Long and Joyner, 1984). For example, Lindsey et al. (1982) reported *Isospora suis* oocysts were unable to sporulate at temperatures above 37°C. Furthermore, the majority of non-gekkonid lizards were observed to be most active during the warmer periods of the day. If lizards defecated at these times, it would serve to limit viability of oocysts that have exogenous sporulation.

We believe the reason that gekkonids have a 10-fold prevalence of infection, when compared to other lizards in this study, is also related to their ecology. Most gekkonids are diverse nocturnal lizards that are distributed throughout the tropics and subtropics of the Old and New Worlds. All 4 species of geckos in this study (see Table 1) are Old World natives that have been introduced into North America and have established themselves in buildings, docks in tropical seaports, in cargoes of fruit, and other humid areas. Feces deposited in these microhabitats would tend to favor increased viability of coccidian oocysts when compared to lizards inhabiting more hostile and dryer sites. Interestingly, these data showing a lower prevalence of infection for coccidia in non-gekkonid lizards are similar to those reported for western rodents inhabiting arid environments with a low prevalence of coccidia (Ford et al., 1990; McAllister et al., 1993).

We thank the Arkansas Game and Fish Commission for Scientific Collecting Permit no. 775,

the New Mexico Department of Game and Fish for permit no. 1804, the Oklahoma Department of Wildlife Conservation for permit no. 1422, and the Texas Parks and Wildlife Department for permit no. SPR-0390-027 (all issued to C.T.M.). We also thank D. M. Boyer, P. S. Freed, E. A. Liner, and J. R. Dixon for providing fecal specimens from some lizards and L. D. Gage for field assistance.

Literature Cited

- Atkinson, C. T., and S. C. Ayala. 1987. *Isospora manchaensis* n. sp., an intranuclear coccidian from the Louisiana ground skink, *Scincella lateralis* (Say, 1823) (Lacertilia: Scincidae). *Journal of Parasitology* 73:817-823.
- Bovee, E. C. 1966. *Eimeria urosaurus* n. sp., a coccidium from the lizard *Urosaurus graciosus* Hallowell, in California. *Journal of Protozoology* 13: 607-610.
- . 1969. *Eimeria umis* n. sp., a coccidian parasite from the lizard *Uma notata*. *Transactions of the American Microscopical Society* 88:257-265.
- , and S. R. Telford, Jr. 1965a. *Eimeria sceloporis* and *Eimeria molochis* spp. n. from lizards. *Journal of Parasitology* 51:85-94.
- , and ———. 1965b. *Eimeria noctisaurus* sp. n., a coccidian from the lizard *Klauberina riversiana*. *Journal of Parasitology* 51:325-330.
- Clark, G. W. 1970. *Eimeria ahtanumensis* n. sp. from the northwestern fence lizard *Sceloporus occidentalis* in central Washington. *Journal of Protozoology* 17:521-530.
- , and D. A. Colwell. 1973. Incidence of eimerians in *Sceloporus occidentalis* from central Washington. *Journal of Parasitology* 59:931-932.
- Collins, J. T. 1990. Standard common and current scientific names for North American amphibians and reptiles, 3rd ed. Society for the Study of Amphibians and Reptiles Herpetological Circular 19: 1-41.
- Conant, R., and J. T. Collins. 1991. *A Field Guide to Reptiles and Amphibians of Eastern and Central North America*. Houghton Mifflin, Boston. 450 pp.
- Ford, P. L., D. W. Duszynski, and C. T. McAllister. 1990. Coccidia (Apicomplexa) from heteromylid rodents in the southwestern United States, Baja California, and northern Mexico with three new species from *Chaetodipus hispidus*. *Journal of Parasitology* 76:325-331.
- Frost, D. R., and R. Etheridge. 1989. A phylogenetic analysis and taxonomy of iguanian lizards (Reptilia: Squamata). University of Kansas Museum of Natural History Miscellaneous Publication 81: 1-65.
- Lindsey, D. S., W. L. Current, and J. V. Ernst. 1982. Sporogony of *Isospora suis* Biester, 1934 of swine. *Journal of Parasitology* 68:861-865.
- Long, P. L., and L. P. Joyner. 1984. Problems in the identification of species of *Eimeria*. *Journal of Protozoology* 31:535-541.
- Matuschka, F.-R. 1989. *Isospora viridanae* n. sp., an intranuclear coccidian parasite from the Canarian skink, *Chalcides viridanus* (Lacertilia: Scincidae). *Journal of Protozoology* 36:274-279.
- , and B. Bannert. 1987. New eimeriid coccidia from the Canarian lizard, *Gallotia galloti* Oudart, 1839. *Journal of Protozoology* 34:231-235.
- McAllister, C. T., and S. J. Upton. 1989. Redescription of *Eimeria boveri* (Apicomplexa: Eimeriidae) from *Hemidactylus mabouia* (Sauria: Gekkonidae), and a new host record for *Eimeria sceloporis*. *Transactions of the American Microscopical Society* 108:92-95.
- , ———, and D. M. Boyer. 1990. *Eimeria dixoni* sp. n. (Apicomplexa: Eimeriidae) from an introduced population of common house geckos, *Hemidactylus frenatus* (Sauria: Gekkonidae), in Dallas County, Texas. *Journal of the Helminthological Society of Washington* 57:1-4.
- , ———, and P. S. Freed. 1988. *Eimeria lineri* sp. n. (Apicomplexa: Eimeriidae) from the Mediterranean gecko, *Hemidactylus turcicus* (Sauria: Gekkonidae), in Louisiana and Texas. *Proceedings of the Helminthological Society of Washington* 55:256-259.
- , ———, R. R. Hollander, and K. M. Hogan. 1993. Coccidian parasites of heteromylid and murid rodents from Baja California del Sur, Mexico. *Journal of the Helminthological Society of Washington* 60:256-258.
- , ———, and S. E. Trauth. 1991a. A new species of coccidia (Apicomplexa: Eimeriidae) from the prairie racerunner, *Cnemidophorus sexlineatus viridis* (Sauria: Teiidae). *Journal of Parasitology* 77:910-913.
- , P. S. Freed, S. J. Upton, D. A. Burdick, and N. A. Wilson. 1991b. Parasites of the rough-tail gecko, *Cryptopodion scabrum* (Sauria: Gekkonidae) from Galveston, Texas. *Texas Journal of Science* 43:199-204.
- Mitschler, R. R., R. L. Morrison, S. R. Goldberg, and S. J. Upton. 1993. Redescription of *Eimeria sceloporis* (Apicomplexa: Eimeriidae) from a new host *Sceloporus jarrovi* (Sauria: Phrynosomatidae). *Texas Journal of Science* 45:356-358.
- Pellérdy, L. P. 1974. *Coccidia and Coccidiosis*, 2nd ed. Verlag Paul Parey, Budapest. 959 pp.
- Telford, S. R., Jr. 1970. A comparative study of endoparasitism among some southern California lizard populations. *American Midland Naturalist* 83: 516-554.
- Upton, S. J., C. T. McAllister, and P. S. Freed. 1988. *Eimeria turcicus* n. sp. (Apicomplexa: Eimeriidae) from the Mediterranean gecko, *Hemidactylus turcicus* (Sauria: Gekkonidae). *Journal of Protozoology* 35:24-25.
- , ———, and S. E. Trauth. 1991. Two new species of coccidia (Apicomplexa: Eimeriidae) from *Eumeces fasciatus* (Sauria: Scincidae) in Arkansas. *Canadian Journal of Zoology* 69:2028-2030.